

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



F.S.A.

FLORASULAM 50

HERBICIDE

ACTIVE CONSTITUENT: 50 g/L FLORASULAM

GROUP	2	HERBICIDE
-------	----------	-----------

A suspension concentrate formulation for post-emergent control of certain broadleaf weeds in cereals and fallow as per the Directions for Use.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT



DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to crops or weeds which may be stressed due to prolonged periods of extreme cold, sustained high temperature within 48 hours of application, moisture stress (water logging or drought), nutritional stress, root disease or previous herbicide treatment, as crop damage or reduced levels of control may result.

DO NOT apply if rain is likely to occur within 3 hours or if heavy rain or storms are forecast within 48 hours.

DO NOT irrigate to the point of run-off for at least 48 hours after application.

DO NOT apply later than growth stage Z37 (flag leaf just visible).

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets are not smaller than a COARSE spray droplet size category
- minimum distances between the application site and downwind sensitive areas are observed. See "Mandatory Downwind Buffer Zones" section of the following table titled "Buffer Zones for Boom Sprayers"
- VERY COARSE when the primary tank mix partner is Amicide Advance 700 or Estercide Xtra 680.

Buffer Zones for Boom Sprayers

Application Rate	Boom height above target canopy	Mandatory Downwind Buffer Zones	
		Natural aquatic areas	Vegetation areas
100 mL/ha	0.5 m or lower	Not required	35 m
	1.0 m or lower	20 m	100 m
100 mL/ha + 440 mL/ha Nufarm Polo® LVE 570 Herbicide	0.5 m or lower	25 m	40 m
	1.0 m or lower	55 m	110 m
100 mL/ha + 500 mL/ha Nufarm Bromicide® MA	0.5 m or lower	25 m	35 m
	1.0 m or lower	55 m	100 m
75 - 100 mL/ha + 260 - 350 mL/ha Nufarm Amicide® Advance 700	0.5 m or lower	30 m	55 m
	1.0 m or lower	70 m	170 m
75 - 100 mL/ha + 265 - 360 mL/ha Nufarm Estericide® Xtra 680	0.5 m or lower	0 m	65 m
	1.0 m or lower	20 m	220 m

DO NOT apply by aircraft unless the following requirements are met:

- spray droplets are not smaller than a COARSE spray droplet size category (VERY COARSE when the primary tank mix partner is Nufarm Amicide® Advance 700 or Nufarm Estericide® Xtra 680)
- for maximum release heights above the target canopy of 3 m or 25% of wingspan or 25% of rotor diameter whichever is the greatest, minimum distances between the application site and downwind sensitive areas are observed. See "Mandatory Downwind Buffer Zones" section of the following table titled "Buffer Zones for Aircraft" -

Buffer Zones for Aircraft

Application Rate	Type of aircraft	Mandatory Downwind Buffer Zones	
		Natural aquatic areas	Vegetation areas
100 mL/ha	Fixed wing	75 m	375 m
	Helicopter	60 m	220 m
100 mL/ha + 440 mL/ha Nufarm Polo® LVE 570 Herbicide	Fixed wing	200 m	425 m
	Helicopter	140 m	240 m
100 mL/ha + 500 mL/ha Nufarm Bromicide® MA	Fixed wing	200 m	375 m
	Helicopter	140 m	230 m
75 - 100 mL/ha + 260 - 350 mL/ha Nufarm Amicide® Advance 700	Fixed wing	210 m	500 m
	Helicopter	150 m	275 m
75 - 100 mL/ha + 265 - 360 mL/ha Nufarm Estericide® Xtra 680	Fixed wing	75 m	600 m
	Helicopter	60 m	300 m

SECTION 1: CROPS AND SITUATIONS

CROP/SITUATION	CROP STAGE	RATE	CRITICAL COMMENTS
<p>F.S.A. Florasulam 500 Herbicide alone or in mixtures with Nufarm Bromicide® MA or Nufarm Polo 570 LVE:</p> <p>Barley Oats Triticale Wheat Ryegrass pasture</p>	3 leaf to flag leaf just visible (Z13-37)	<p>100 mL/ha + 0.5 L/100 L Nufarm CanDo™ Adjuvant*</p> <p>OR</p> <p>100 mL/ha + 500 mL/ha Nufarm Bromicide® MA + 0.5 L/100 L Nufarm CanDo™ Adjuvant*</p> <p>OR</p> <p>100 mL/ha + 440 mL/ha Nufarm Polo 570 LVE + 0.5 L/100 L Nufarm CanDo™ Adjuvant*</p>	<p>F.S.A. Florasulam 50 Herbicide will not reliably control weeds that emerge after treatment. Apply to actively growing weeds. Best results are achieved when weeds are small (2 - 4 leaf) and under good growing conditions. Treatment of crop or weeds that are stressed must be avoided. While grain yields are unaffected, slight, transient yellowing and growth retardation may occur in some conditions. Crop recovery is usually rapid, but may be slower if crops are stressed due to prolonged periods of extreme cold, moisture stress, nutritional stress or root disease. F.S.A. Florasulam 50 Herbicide can be tank mixed with compatible broad-leaf herbicides to extend the weed spectrum as per the labels of those products.</p> <p><u>In fallows:</u> To add control of grass weeds, F.S.A. Florasulam 50 Herbicide (including in mixtures with Nufarm Amicide Advance® 700 or Nufarm Estercide Xtra® 680), can be mixed with Nufarm CRUCIAL®**. Refer to the Nufarm CRUCIAL® label for use rates and adjuvants recommended for the grasses (see General Instructions, Compatibility Section).</p>
<p>F.S.A. Florasulam 50 Herbicide in mixtures with Nufarm Amicide® Advance 700:</p> <p>Barley Oats Triticale Wheat Ryegrass pasture</p>	5 leaf to flag leaf just visible (Z15-37)	<p>75 - 100 mL/ha + 260 - 350 mL/ha Nufarm Amicide Advance® 700 + 0.5 L/100 L Nufarm CanDo™ Adjuvant*</p>	
<p>F.S.A. Florasulam 50 Herbicide alone or in mixtures with Nufarm Amicide® Advance 700 or Nufarm Estercide® Xtra 680:</p> <p>Summer or winter fallow prior to planting crops listed in the ROTATIONAL CROPS section.</p>	Fallow	<p>100 mL/ha + 0.5 L/100 L Nufarm CanDo™ Adjuvant*</p> <p>OR</p> <p>75 - 100 mL/ha + 260 - 350 mL/ha Nufarm Amicide Advance® 700 + 0.5 L/100 L Nufarm CanDo™ Adjuvant*</p> <p>OR</p> <p>75 - 100 mL/ha + 265 - 360 mL/ha Nufarm Estercide Xtra® 680 + 0.5 L/100 L Nufarm CanDo™ Adjuvant*</p>	

** Alternatively to Nufarm CRUCIAL®, Nufarm Weedmaster DST, Nufarm Weedmaster ARGO or Nufarm Gladiator® CT can be used provided all label guidelines are observed.

* Alternatively to Nufarm CanDo™ Adjuvant, Banjo® or another registered high quality methylated seed oil adjuvant can be used at 1 L/100L.

SECTION 2: WEED TABLE

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE F.S.A. Florasulam 50 Herbicide used alone	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Bromicide® MA	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Polo® 570 LVE	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Amicide® Advance 700	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Estericide® Xtra 680	CRITICAL COMMENTS
		ALWAYS add 0.5 L/100 L Nufarm CanDo™ Adjuvant					
Bedstraw (<i>Galium tricornutum</i>)	Up to 4 whorl or 8 cm in height	100 mL/ha F.S.A. Florasulam 50 Herbicide	100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			Addition of Nufarm Bromicide® MA or Nufarm Polo® 570 LVE adds improved early brownout.
	Up to 6 whorl or 10 cm in height				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estericide® Xtra 680	
Bifora (<i>Bifora testiculata</i>)	Up to 4 leaf or 5 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide					Control of Bifora can be expected if actively growing weeds are treated in warm conditions. Conversely, control may be reduced in cold, wet conditions that slow plant growth.
Capeweed (<i>Arctotheca calendula</i>)	Up to 6 leaf or 8 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide	100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA				Control of Capeweed can be expected if actively growing weeds are treated in warm conditions. Conversely, control may be reduced in cold, wet conditions that slow plant growth. Addition of Nufarm Bromicide® MA adds improved final control.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE F.S.A. Florasulam 50 Herbicide used alone	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Bromicide® MA	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Polo® 570 LVE	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Amicide® Advance 700	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Estercide® Xtra 680	CRITICAL COMMENTS
		ALWAYS add 0.5 L/100 L Nufarm CanDo™ Adjuvant					
Charlock (<i>Sinapis arvensis</i>)	Up to 4 leaf		100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			
Climbing buckwheat/black bindweed (<i>Fallopia convolvulus</i>)	Up to 6 leaf				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide® Xtra 680	Suppression only.
					75 mL/ha F.S.A. Florasulam 50 Herbicide + 260 mL/ha Nufarm Amicide Advance® 700 + Trooper® 242 at label rates		The addition of Trooper® 242 adds control of Climbing buckwheat/black bindweed in NSW and Qld only with a lower rate of F.S.A. Florasulam 50 Herbicide and Nufarm Amicide Advance® 700. DO NOT add Trooper® 242 when applying in fallows.
Common Fumitory (<i>Fumaria officinalis</i>)	Up to 4 leaf or 5 cm in height	100 mL/ha F.S.A. Florasulam 50 Herbicide					Suppression only.
Dense-flowered Fumitory (<i>Fumaria densiflora</i>)	Up to 3 leaf or 4 cm in height	100 mL/ha F.S.A. Florasulam 50 Herbicide		100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo 570 LVE			Addition of Nufarm Polo® 570 LVE adds improved final control.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE F.S.A. Florasulam 50 Herbicide used alone	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Bromicide® MA	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Polo® 570 LVE	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Amicide® Advance 700	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Estercide® Xtra 680	CRITICAL COMMENTS
		ALWAYS add 0.5 L/100 L Nufarm CanDo™ Adjuvant					
Double-gee/Spiny Emex/Three Cornered Jack (<i>Emex australis</i>)	Up to 4 leaf or 10 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide	100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA		100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide® Xtra 680	Addition of Nufarm Bromicide® MA adds improved early brownout.
Hedge Mustard (<i>Sisymbrium officinale</i>)	Up to 4 leaf	100 mL/ha F.S.A. Florasulam 50 Herbicide	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			Addition of Nufarm Bromicide® MA or Nufarm Polo® 570 LVE adds improved early brownout.
Indian hedge mustard (<i>Sisymbrium orientale</i>)	Up to 4 leaf				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide® Xtra 680	
	Up to 6 leaf		100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			
Long Storksbill (<i>Erodium botrys</i>)	Up to 9 leaf	100 mL/ha F.S.A. Florasulam 50 Herbicide					
Matricaria (<i>Oncosiphon piluliferum</i>)	Up to 20 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide					Suppression only.
Mexican Poppy (<i>Argemone mexicana</i>)	Up to 4 leaf or 4 cm in height		100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA				

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE F.S.A. Florasulam 50 Herbicide used alone	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Bromicide® MA	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Polo® 570 LVE	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Amicide® Advance 700	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Estercide® Xtra 680	CRITICAL COMMENTS
		ALWAYS add 0.5 L/100 L Nufarm CanDo™ Adjuvant					
Prickly Lettuce (<i>Lactuca serriola</i>)	Up to 7 leaf or 6 cm in diameter		100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA				
Shepherd's Purse (<i>Capsella bursa-pastoris</i>)	Up to 8 leaf or 5 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide	100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			Addition of Nufarm Bromicide® MA or Nufarm Polo® 570 LVE adds improved early brownout.
Snail Medic (<i>Medicago scutellata</i>)	Up to 6 leaf or 10 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide	100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			Addition of Nufarm Bromicide® MA or Nufarm Polo® 570 LVE adds improved early brownout.
Sowthistle/ Milk Thistle (<i>Sonchus oleraceus</i>)	Up to 3 leaf or 2 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide					
Sub-clover (<i>Trifolium subterraneum</i>)	Up to 6 leaf or 10 cm in diameter	100 mL/ha F.S.A. Florasulam 50 Herbicide					
Turnip weed (<i>Rapistrum rugosum</i>)	Up to 4 leaf				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide® Xtra 680	
	Up to 6 leaf		100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE F.S.A. Florasulam 50 Herbicide used alone	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Bromicide® MA	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Polo® 570 LVE	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Amicide® Advance 700	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Estercide® Xtra 680	CRITICAL COMMENTS
ALWAYS add 0.5 L/100 L Nufarm CanDo™ Adjuvant							
Variegated thistle (<i>Silybum marianum</i>)	Up to 6 leaf				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide® Xtra 680	Suppression only.
					75 mL/ha F.S.A. Florasulam 50 Herbicide + 260 mL/ha Nufarm Amicide Advance® 700 + Trooper® 242 at label rates		The addition of Trooper® 242 adds control of Variegated thistle in NSW and Qld only with a lower rate of F.S.A. Florasulam 50 Herbicide and Nufarm Amicide Advance® 700. DO NOT add Trooper® 242 when applying in fallows.
Vetch/Tares (<i>Vicia sativa</i>)	Up to 5 node or 10 cm in height	100 mL/ha F.S.A. Florasulam 50 Herbicide	100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			Addition of Nufarm Bromicide® MA or Nufarm Polo 570 LVE adds improved early brownout.
	Up to 6 node or 20 cm in height				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide Xtra® 680	
Volunteer Canola (<i>Brassica napus</i>) (non-imidazolinone herbicide tolerant varieties)	Up to 6 leaf	100 mL/ha F.S.A. Florasulam 50 Herbicide					

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE F.S.A. Florasulam 50 Herbicide used alone	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Bromicide® MA	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Polo® 570 LVE	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Amicide® Advance 700	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Estercide® Xtra 680	CRITICAL COMMENTS
		ALWAYS add 0.5 L/100 L Nufarm CanDo™ Adjuvant					
Volunteer Chickpea (<i>Cicer arietinum</i>)	Up to 5 node	100 mL/ha F.S.A. Florasulam 50 Herbicide					
	Up to 6 node or 20 cm in height				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide Xtra® 680	
Volunteer Faba Bean (<i>Vicia faba</i>) (non-imidazolinone herbicide tolerant varieties)	Up to 3 node	100 mL/ha Nufarm Saracen®	100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			Addition of Nufarm Bromicide® MA or Nufarm Polo® 570 LVE adds improved early brownout.
	Up to 6 node or 15 cm in height				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide Xtra® 680	
Volunteer Field Pea (<i>Pisum sativum</i>) (non-imidazolinone herbicide tolerant varieties)	Up to 6 node or 12 cm in height	100 mL/ha F.S.A. Florasulam 50 Herbicide					
Volunteer lentil (<i>Lens culinaris</i>)	Up to 6 node or 20 cm in height				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide Xtra® 680	

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE F.S.A. Florasulam 50 Herbicide used alone	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Bromicide® MA	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Polo® 570 LVE	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Amicide® Advance 700	RATE F.S.A. Florasulam 50 Herbicide + Nufarm Estercide® Xtra 680	CRITICAL COMMENTS
		ALWAYS add 0.5 L/100 L Nufarm CanDo™ Adjuvant					
Volunteer Lupins (<i>Lupinus angustifolius</i>)	Up to 6 leaf or 15 cm in height	100 mL/ha F.S.A. Florasulam 50 Herbicide®		100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			Addition Nufarm Polo® 570 LVE adds improved early brownout.
Wild radish (<i>Raphanus raphanistrum</i>)	Up to 4 leaf or 5 cm in diameter		100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE	100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide Xtra® 680	
Wild turnip (<i>Brassica tournefortii</i>)	Up to 4 leaf				100 mL/ha F.S.A. Florasulam 50 Herbicide + 350 mL/ha Nufarm Amicide Advance® 700	100 mL/ha F.S.A. Florasulam 50 Herbicide + 360 mL/ha Nufarm Estercide Xtra® 680	
	Up to 5 leaf or 10 cm in diameter		100 mL/ha F.S.A. Florasulam 50 Herbicide + 500 mL/ha Nufarm Bromicide® MA	100 mL/ha F.S.A. Florasulam 50 Herbicide + 440 mL/ha Nufarm Polo® 570 LVE			

* Alternatively to Nufarm CanDo™ Adjuvant, Banjo® or another registered high quality methylated seed oil adjuvant can be used at 1 L/100L.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

When using F.S.A. Florasulam 50 in tank mix with another product, observe the Harvest Withholding Period of the other product.

GRAZING: DO NOT CUT OR GRAZE FOR STOCKFOOD FOR 7 DAYS AFTER APPLICATION

When using F.S.A. Florasulam 50 in tank mix with another product, observe which ever grazing Withholding Period is longer.

TRADE ADVICE

FODDER INTENDED FOR EXPORT: When F.S.A. Florasulam 50 Herbicide is used as directed residues in fodder are not expected. However, as export requirements are subject to change, consult your exporter before using this product on crops destined to be used for export fodder.

LIVESTOCK DESTINED FOR EXPORT MARKETS: When F.S.A. Florasulam 50 Herbicide is used as directed and the above withholding period is observed, livestock commodities are considered acceptable for export. However, as export requirements are subject to change, consult your exporter for updated information about specific market requirements.

GENERAL INSTRUCTIONS

F.S.A. Florasulam 50 Herbicide is a suspension concentrate formulation that mixes readily with water. F.S.A. Florasulam 50 Herbicide is a foliar herbicide for post-emergence use in wheat, barley, triticale, oats and ryegrass pasture. It will not reliably control weeds that emerge after treatment. Apply to actively growing weeds. Best results are achieved when weeds are small (2 - 4 leaf) and under good growing conditions. Treatment of crop or weeds that are stressed must be avoided.

Crop Safety

While grain yields are unaffected, slight, transient yellowing and growth retardation may occur in some conditions. Crop recovery is usually rapid, but may be slower if crops are stressed due to prolonged periods of extreme cold, moisture stress, nutritional stress or root disease. F.S.A. Florasulam 50 Herbicide can be tank mixed with compatible broad-leaf herbicides to extend the weed spectrum as per the labels of those products.

Rotational Crops

F.S.A. Florasulam 50 Herbicide is primarily broken down in soil by microbial activity. Under conditions not conducive to microbial activity, e.g. dry seasons, cold and/or waterlogged soil it may take longer to degrade.

Plantback intervals to winter crops after an application of F.S.A. Florasulam 50 Herbicide

Crop to be sown	Minimum re-cropping interval	Minimum rainfall requirements from application to planting
Wheat, barley, triticale	1 week	None
Oats	6 weeks	25 mm
Sub clover, canola, chickpeas, faba bean, field pea, lentils, lupins, medic, vetch	8 months	100 mm

Plantback intervals to summer crops after an application of F.S.A. Florasulam 50 Herbicide

Crop to be sown	Minimum re-cropping interval	Minimum rainfall requirements from application to planting
Maize, sorghum	4 months	100 mm
Mung beans	5 months	150 mm
All other summer crops including cotton, sunflowers, soybeans	6 months	150 mm

Mixing

Add the required amount of F.S.A. Florasulam 50 Herbicide to the spray tank when partly filled with water and begin agitation. Ensure agitation is maintained while topping up the tank and during application. Add Nufarm CanDo™ Adjuvant at 0.5 L/100 L after F.S.A. Florasulam 50 Herbicide is well mixed in the spray solution and near the end of the filling process to minimize foaming. Only mix sufficient chemical for each day's work. **Tank mixtures:** wettable powder or dry flowable formulations should be added to the spray tank first, followed by suspension concentrates (F.S.A. Florasulam 50 Herbicide), aqueous concentrates (e.g. Nufarm Amicide Advance® 700, Nufarm CRUCIAL®) and then emulsifiable concentrate formulations (e.g. Nufarm Bromicide® MA, Nufarm Polo® 570 LVE, Nufarm Estericide Xtra® 680). **DO NOT** allow prepared mixture to stand overnight.

Boom Sprayer Application

Apply F.S.A. Florasulam 50 Herbicide in sufficient water to obtain good coverage. Apply by an accurately calibrated ground rig using a water volume of 80 - 100 L/ha.

Aerial Application

Apply a minimum water volume of 40 L/ha. Markers should always move upwind before aircraft passes overhead. Application by micronair equipment may give variable results.

Cleaning Spray Equipment

After using F.S.A. Florasulam 50 Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Drain the tank and wash tank, pump, line and nozzle filters.

Partial cleaning (rinse only - before using rig to spray barley and wheat): after cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

Complete cleaning (decontamination - before using rig to spray crops that are susceptible to F.S.A. Florasulam 50 Herbicide): after cleaning the tank as above, quarter fill the tank with clean water and add a liquid alkali detergent at 500 mL/100L water or tank cleaner (e.g. Nufarm Tank and Equipment Cleaner) at recommended rates and circulate throughout the system for at least 15 minutes. Drain the whole system. Then remove filters and nozzles and clean separately. Finally, rinse inside the tank thoroughly using a pressure hose and flush system with clean water and allow to drain.

Note: if using a concentrated laundry detergent use 250 g (or mL)/100L. **DO NOT** use chlorine based cleaners.

Rinse water should be discharged into a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and their roots and watercourses.

COMPATIBILITY

F.S.A. Florasulam 50 Herbicide is compatible with Nufarm Amicide Advance® 700, Nufarm Bromicide® 200, Nufarm Bromicide® MA, Nufarm CRUCIAL® Advanced Technology Herbicide, chlorpyrifos, Nufarm Estericide® Xtra 680, Frequency*, Nufarm Gladiator® CT, Nufarm Kaiso® 240EG, Opus* 125, Pirimor* WG, Nufarm Polo® 570 LVE, Prosaro* 420 SC, Talinor*, Tazer Xpert*, Throttle® 500, Trojan*, Trooper® 242, Velocity*, Nufarm weedmaster ARGO®, Nufarm weedmaster DST®. **Note:** all mix partners must be able to be used with a crop oil. F.S.A. Florasulam 50 Herbicide® has not been tested in all tank mix combinations over all cereal varieties. All mixtures should be tested prior to mixing commercial quantities.

Adjuvants: always use Nufarm CanDo™ Adjuvant (0.5 L/100 L), Banjo® Spray Adjuvant (1 L/100 L) or other registered high quality methylated seed oil adjuvant at the recommended rate.

RESISTANT WEEDS WARNING

GROUP	2	HERBICIDE
-------	----------	-----------

F.S.A. Florasulam 50 Herbicide is a member of the triazolopyrimidine sulfonanilide group of herbicides. F.S.A. Florasulam 50 has the acetolactate synthase (ALS) inhibitor mode of action. For herbicide resistance management, Florasulam 50 is a Group 2 herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 2 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 2 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Four Seasons Agribusiness accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Four Seasons Agribusiness representative.

PRECAUTIONS

RE-ENTRY: DO NOT allow entry into treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When using together with other products, consult their label safety directions. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and goggles. Wash hands after use. After each day's use, wash contaminated clothing, gloves and goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

CONDITIONS OF SALE

The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

DO NOT eat, drink or smoke when using this product. Wear protective gloves, clothing, eye and face protection. Wash hands and exposed skin thoroughly after handling. IF SWALLOWED call the Poison Centre of a doctor if you feel unwell.

**IN AN EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**



Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, New South Wales, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255
www.fourseasonsag.com

IN AN EMERGENCY, DIAL 000, POLICE OR FIRE BRIGADE



Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, New South Wales, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255
www.fourseasonsag.com