

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



F.S.A.

CLOPYRALID 600 HERBICIDE

**ACTIVE CONSTITUENT: 600 g/L CLOPYRALID
present as the acid and TRIISOPROPANOLAMINE SALT**

GROUP

4

HERBICIDE

For the control of a wide range of Broadleaf weeds in wheat, barley, triticale, oats, pastures, canola, fallow land, forests and industrial situations as specified in the directions for use table.

**IMPORTANT: READ THE ATTACHED BOOKLET
BEFORE USING THIS PRODUCT**



DIRECTIONS FOR USE

IT IS ESSENTIAL to select a rate appropriate to weed size. Best results will be obtained when weeds are actively growing at treatment.

Restrictions:

CROP AND FALLOW LAND

RESTRAINTS

DO NOT apply to weeds, which may be stressed (inactive growth) due to prolonged periods of extreme heat or cold, moisture stress (water logging or drought) or previous herbicide treatment as reduced levels of control may result.

DO NOT spray if rain is likely within 3 hours.

DO NOT apply immediately before sowing susceptible crops, or sow susceptible crops into paddocks treated the previous year with F.S.A. Clopyralid 600 Herbicide until after the required plantback period has elapsed (see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section).

DO NOT compost material from treated plants or crops before reading the PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.

FORESTS

RESTRAINTS

DO NOT apply to weeds which may be stressed (inactive growth) due to prolonged periods of extreme heat or cold, moisture stress (waterlogging or drought) or previous herbicide treatment as reduced levels of control may result.

DO NOT spray if rain is likely.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

1. CROPS, PASTURES AND FALLOW LAND

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Barley Oats Triticale Wheat	Pre-sowing	Capeweed	Up to 8 leaf and maximum 10 cm diameter	75 plus knockdown herbicide	WA only	Pre-sowing: This rate should only be used in tank mixture with formulations of paraquat/diquat or glyphosate.
	Post-sowing pre-emergence through to 3 leaf			75 plus diuron at labeled rates		Post sowing pre-emergent to 3 leaf: This rate should only be used in tank mixture with diuron for control of transplants. Refer to diuron label for specific rates.
			Early post-emergence (2 leaf to jointing)	Cotyledons to 6 leaf and maximum 5 cm diameter		75
	4 to 5 leaf onwards	Capeweed, Soldier thistle	Up to 10 cm diameter (4 to 8 leaf)	150	NSW, ACT, Vic, SA, Tas and WA only	Weeds should be young and actively growing. Weeds will become stunted and non-competitive soon after application although final results may not show for some weeks. Faba beans and lupins will only be suppressed.
		Volunteer chickpeas, lentils and safflower	Up to 6 leaf	125		
		Volunteer faba beans and lupins	Up to 4 leaf			
		Volunteer field peas	Maximum 10 cm high or 6 nodes	75 40 plus 700 mL/ha MCPA LVE		
		Volunteer medics and seedling lucerne	Up to 8 leaf	75		
		Volunteer sub-clover	Up to 6 leaf			
		Volunteer vetch	Runners up to 10 cm maximum 16 leaf	50 40 plus 700mL/ha MCPA LVE		
Barley Oats Triticale Wheat	4 to 5 leaf onwards	Prickly lettuce	4 to 6 leaf and maximum 8cm diameter	75 plus 700mL/ha MCPA LVE	NSW, ACT, Vic, SA, Tas and WA only	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks.
		Thistles including: Nodding, Saffron, Scotch, Slender, Spear, Stemless, Variegated	Rosettes up to 10cm maximum diameter	25 plus 1.0L/ha MCPA amine (500g/L) or 25 plus 700mL/ha MCPA LVE		NSW, ACT, Vic, SA, Tas, Qld and WA only
	5 leaf to late tillering	Skeleton weed	5 to 15cm rosettes	250 plus 1.0L/ha MCPA amine (500 g/L)	NSW, ACT, Vic and SA only	Weeds should be a minimum 5cm in diameter and growing actively. This rate will give control until harvest and substantially reduce weed numbers the following season.
Canola	2 to 8 leaf	Capeweed, Saffron thistle, Skeleton weed, Soldier thistle	Up to 10cm diameter (4 to 8 leaf)	150	NSW, ACT, Vic, SA, Tas, WA, Qld only	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks. Skeleton weed will only be controlled until harvest.
		Volunteer chickpeas, lentils and safflower	Up to 6 leaf	125		Faba beans and lupins will only be suppressed.
		Volunteer faba beans and lupins	Up to 4 leaf	125		For the control of annual grasses, F.S.A. Clopyralid 600 Herbicide may be tank mixed with haloxyfop herbicides.
		Volunteer field peas	Maximum 10 cm high or 6 nodes	75		
		Volunteer medics and seedling lucerne	Up to 8 leaf	75		
		Volunteer sub-clover	Up to 6 leaf	75		
		Volunteer vetch	Runners up to 10 cm maximum 16 leaf	50		
Pastures and fallow land	Post-emergence	Hardhead thistle (Creeping knapweed, Russian knapweed)	Actively growing plants	Hand gun: 250mL/100L of water	Vic only	NOTE: DO NOT USE ON LUCERNE. CLOVERS AND MEDICS WILL BE ELIMINATED FOR AT LEAST ONE YEAR. Victoria only: Use the lower rate only on light soils (sand and sandy loam) where a slightly lower degree of control is acceptable. Use the higher rate on all soil types where complete control is required. Addition of a wetting agent at label rates is recommended for treatment of hardhead thistle. Spray between September and April on actively growing plants for effective control. Thorough coverage is essential. Apply in 200 to 250L of water/ha. BOOM SPRAYING: Use the higher rates of F.S.A. Clopyralid 600 Herbicide plus MCPA on multi-crowned plants or rosettes larger than 30cm in diameter. Spraying may be done at any time during active growth, usually in early winter or spring. Avoid spraying during the dormant winter period or at any time when thistles are not actively growing. <i>Do not spray flowering thistles.</i> PRE-SPRAY MANAGEMENT: The pasture should be slightly grazed prior to spraying to reduce clover and grass cover and expose the smaller thistles to the spray. The grazed pasture should be left for seven days to allow thistles to freshen prior to treatment. POST-TREATMENT MANAGEMENT: Response of thistles to treatment with the F.S.A. Clopyralid 600 Herbicide plus MCPA mixture will be slow compared to the standard treatments with 2,4-D or MCPA. If possible, delay grazing of sprayed thistles for 14 days after treatment. CLOVER DAMAGE: The F.S.A. Clopyralid 600 Herbicide plus MCPA mixture can be very damaging to subterranean clover. The lower rate is no more damaging than label rates of 2,4-D or MCPA. The higher rate of the F.S.A. Clopyralid 600 Herbicide plus MCPA mixture will reduce the clover component of the pasture for about two months. Clover recovery will be quicker during periods of active growth. HANDGUN (Spot spray): Treat from rosette stage to early flowering. Thorough spraying is necessary. DRENCHGUN: Apply 5 mL to rosette crown. To multicrown plants, apply 5 mL to each crown.
				Boom spray: 1 or 2L/ha		
		Thistles including: Nodding, Variegated, Scotch, Spear, Slender, Saffron	Rosette stage prior to stem elongation. Treat rosette stage prior to stem elongation.	25 or 35 mL/ha plus 1 to 1.5L MCPA amine (500g/L)/ha Drench gun: 25 mL/1L of water Hand gun: 125 mL/100L of water	NSW, ACT, Vic, Tas, SA and Qld only	
			Nodding thistle	Rosettes up to 20cm diameter	50	
Californian thistle	From early buds to flowering (December to February)	Hand gun: 125 mL/100L of water Boom spray: 1 L/ha	Vic and Tas only	Addition of a wetting agent at label rates is recommended. Re-treatment of re-growth in the year following treatment will usually be necessary to achieve a high level of control. NOTE: Clovers and medics will be eliminated for at least one year.		
Pasture, Forests, Rights of-Way, Industrial Situations	Post-emergence	Groundsel bush	Young seedlings to mature plants	Hand gun: 165 or 250 mL/100L of water	Qld, NSW, ACT only	Spray foliage when growth is active. Use the lower rate on young seedlings and the higher rate on plants more than 2m tall or when growth is slow.
Pasture	Post-emergence	Silver wattle	Active growth spring to summer	Hand gun: 250 mL/ 100L of water Boom and aerial spray: 2.5 L/ha (weeds 30cm to 2m) 3.5 L/ha (weeds 2 to 4m) 4.25 L/ha (weeds 4 to 8m)	NSW, ACT, Vic and Tas only	For effective control apply when bushes are growing actively. Large trees will not show complete necrosis. HANDGUN: Means high volume NOT low volume knapsack. Spray to give full coverage of leaves and stems. Add organosilicone surfactant at 200 mL/100L for optimum results. See General Instructions for handgun spraying. BOOM AND AERIAL SPRAYING: For boom spraying apply in 150 to 200L of water/ha. For aerial treatment apply in a minimum of 50L/ha of water containing 25 to 50% by volume of anti-evaporant oil such as Ulvapron. Mix F.S.A. Clopyralid 600 Herbicide and water first and then add Ulvapron. Maintain continuous agitation. Clovers and other legumes will be eliminated for at least one year.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

2. FORESTS

DIRECTIONS FOR USE: It is essential to select a rate appropriate to weed size. Best results will be obtained when weeds are actively growing at treatment.

Pre-planting: Boom and Aerial Application.

IMPORTANT NOTE: For aerial application on plantation trees including *Pinus radiata* plantations, a closed mixing and loading system MUST be used.

FORESTS AND PLANTATION TREES INCLUDING <i>EUCALYPTUS</i> SPP., <i>CORYMBIA MACULATA</i> AND <i>PINUS RADIATA</i>				
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	STATE	CRITICAL COMMENTS
Capeweed, Thistles, Volunteer legumes, Flatweed, Fleabanes	Pre-emergent	1-3 L	All States	Use the higher rate for extended pre-emergence control (>3 months).
<i>PINUS RADIATA</i> only				
Silver Wattle (suppression only)	Pre-emergence from seeds	3 L	NSW, ACT, Vic, Tas, SA only	For best results apply F.S.A. Clopyralid 600 to bare soil just prior to Spring rain or when wattles are expected to germinate. Avoid application to heavy trash situations. A high level of suppression may not be achieved where rain does not fall for an extended period after application (>1 month), or where very high rainfall occurs after application (>1200 mm/yr).

Post-Planting: High Volume Spraying by Handgun

FORESTS AND PLANTATION TREES INCLUDING <i>EUCALYPTUS</i> SPP., <i>CORYMBIA MACULATA</i> AND <i>PINUS RADIATA</i>				
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /100 L WATER	STATE	CRITICAL COMMENTS
Groundsel Bush	Young seedlings to mature plants	165 – 250 mL	Qld, NSW, ACT only	Do not apply by using spraying equipment carried on the back of the user. Spray foliage when growth is active. Use the lower rate on young seedlings and the higher rate on plants more than 2 m tall or when growth is slow.
Ragwort	Actively growing rosettes up to stem elongation and before flowering	100 to 150 mL	All States	Do not apply by using spraying equipment carried on the back of the user. Spray from the rosette to the shooting stage of growth. Use the higher rate on large multi-crown plants. Addition of a 100% non-ionic surfactant such as WetDrop Wetter at 0.1% v/v is recommended. Add Diquat (200 g/L) at 1 L/100 L water plus a surfactant after opening of the first flowers, to prevent the formation of viable seed. Where Diquat is added use a directed spray to avoid tree injury.
Silver Wattle	Active growth Spring to Summer	250 mL	NSW, ACT, Vic, Tas, SA only	For effective control apply when bushes are growing actively. Large trees will not show complete necrosis. Handgun: Means high volume NOT low volume knapsack. Do not apply by using spraying equipment carried on the back of the user. (See General Instructions, Application). Spray to the point of run-off to give full coverage of leaves and stems. Add Pulse (organosilicone surfactant) at 200 mL/100 L for optimum results.
Cape Ivy	Any Growth Stage	1.65 L	Vic, Tas only	Do not apply by using spraying equipment carried on the back of the user. Apply by handheld weed wiper or CDA at dilutions with water at 1:3. Application may be made at any time of the year provided foliage is dry at the time. Avoid spraying non-target plants.

Post-Planting: Boom and Aerial Application

IMPORTANT NOTE: For aerial application on plantation trees including *Pinus radiata* plantations, a closed mixing and loading system **MUST** be used.

FORESTS AND PLANTATION TREES INCLUDING <i>EUCALYPTUS</i> SPP., <i>CORYMBIA MACULATA</i> AND <i>PINUS RADIATA</i>				
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	STATE	CRITICAL COMMENTS
Capeweed, Flatweed, Thistles (except Hardhead Thistle), Volunteer Legumes, Skeleton Weed	Actively growing rosettes, seedlings up to 15 cm diameter or height.	250 to 500 mL	All States	Cupping of the tip leaves and "weepy leader" symptoms may occur on certain Eucalyptus spp. and Corymbia maculata and are generally transient and do not result in long-term injury. These symptoms may be more obvious at rates of 1 L/ha or higher or where mixtures are used on blue gum, shining gum and spotted gum. Where "weepy leader" effect is a concern use a directed spray. Use the 250 mL rate until 3 months post planting and the 500 mL rate for trees 3 months and older. Use the low rate only under ideal conditions with excellent weed growth and where knockdown control of small weeds is desired. Use the high rate where longer control is required of larger weeds. For the control of annual and certain perennial grasses, F.S.A. Clopyralid 600 can be tank mixed with Verdict* or other 520 g/L Haloxyfop products. See also comments in Mixing section. Bonza* should not be used in tank mixes with Verdict or other 520 g/L Haloxyfop products and F.S.A. Clopyralid 600 on sensitive species such as blue gum, shining gum or spotted gum where rates of F.S.A. Clopyralid 600 are more than 1 L/ha. Use a 100% non-ionic surfactant such as WetDrop Wetter at 0.1% v/v instead.
Capeweed, Flatweed, Fleabane, Thistles including Hardhead Thistle, Volunteer Legumes, Skeleton Weed	Actively growing rosettes and seedlings greater than 15 cm diameter or height up to stem elongation and before flowering	1 L		
Californian Thistle	From early bud to flowering (December to February)			For best control of California Thistle use a wetter such as WetDrop Wetter at 0.1% v/v. A second annual application may also be required for the best control.
Ragwort	Small rosettes to larger rosettes up to stem elongation and before flowering	500 mL to 1 L		Spray from the rosette to the shooting stage of growth. For small rosette seedling plants use the lower rate. For large rosette multicrown and/or perennial plants use the higher rate. Addition of a 100% non-ionic surfactant such as WetDrop Wetter at 0.1% v/v is recommended. Add Diquat (200 g/L) at 1 L/100L water plus a surfactant after opening of the first flowers, to prevent the formation of viable seed. Where diquat is added use a directed spray to avoid tree injury.

FORESTS AND PLANTATION TREES INCLUDING <i>EUCALYPTUS</i> SPP., <i>CORYMBIA MACULATA</i> AND <i>PINUS RADIATA</i>				
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	STATE	CRITICAL COMMENTS
Sorrel (suppression only)	Actively growing rosettes, seedlings up to 15 cm diameter or height	3 to 4.25 L	All States	Higher rates give better suppression. At rates greater than 3 L use a directed spray to avoid tree injury.
Silver Wattle	Active growth Spring to Summer (0.5 to 2 m tall)	2.5 L	NSW, ACT, Vic, Tas, SA only	For effective control apply when bushes are growing actively. Large trees will not show complete necrosis. For boom spraying apply in 150 to 200 water/ha. For aerial treatment apply in a minimum of 50 L/ha of water containing 25 to 50% by volume of anti-evaporant such as Broadcoat*. Mix F.S.A. Clopyralid 600 and water first and then add Broadcoat*. Maintain continuous agitation. At rates of 3.5 L and 4.25 L for Eucalypt spp. use a directed spray to avoid tree injury.
	Active growth Spring to Summer (2 to 4 m tall)	3.5 L		
	Active growth Spring to Summer (4 to 8 m tall)	4.25 L		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Harvest

CEREALS: DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

CANOLA: DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION.

FOREST: NOT REQUIRED WHEN USED AS DIRECTED.

Grazing

CEREALS, CANOLA, PASTURES and FALLOW LAND: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

FORESTS, except *Pinus radiata* PLANTATIONS: DO NOT GRAZE FOR 7 DAYS AFTER APPLICATION.

***Pinus radiata* PLANTATIONS: DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION.**

GENERAL INSTRUCTIONS

MIXING

Half fill the spray tank with water and add the required amount of F.S.A. Clopyralid 600 Herbicide and complete filling. Agitate continuously to ensure thorough mixing before and during application. Only mix sufficient chemical for each day's work.

Aerial Application – Plantation trees including *Pinus radiata* plantations: Closed mixing and loading system MUST be used.

Tank mixtures

Wettable powder or dry flowable formulations should be added to the spray tank first, followed by suspension concentrates (flowables), aqueous concentrates, emulsifiable concentrate formulations (e.g. Haloxypop or MCPA LVE).

COMPATIBILITY

F.S.A. Clopyralid 600 Herbicide is compatible with the following:

BROADLEAF HERBICIDES: Metsulfuron-methyl, bromoxynil, chlorsulfuron, diuron, glyphosate, MCPA amine, MCPA LVE, paraquat, paraquat/diquat, terbutryn, 2,4-D amine.

GRASS HERBICIDES IN CEREAL CROPS: diclofop-methyl, Grasp +, Puma + (+ Grasp and Puma for wild oat control only).

GRASS HERBICIDES ON BROADLEAF CROPS: Haloxyfop 520

APPLICATION

BOOM SPRAYING CROP and PASTURES

Apply F.S.A. Clopyralid 600 Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft, delivering 200 to 300 micron droplets and not less than 50L/ha water volume for boom sprayers or not less than 20L/ha for aerial applications.

Hardhead thistle - Use a spray volume of 200 to 250L/ha of water.

Silver wattle - Use a spray volume of 150 to 200L/ha of water by ground boomspray and a minimum spray volume of 50L/ha by aircraft.

Boom Spraying Plantation Trees

Apply F.S.A. Clopyralid 600 Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft, delivery Medium to Coarse droplets size and not less than 50 L/ha water volume for boom sprayers or not less than 20 L/ha for aerial application.

HIGH VOLUME HANDGUN

Apply the recommended mix to give full coverage of leaves and stems through a No. 6-8 tip at 700 to 1500 kPa. Spray volume for effective coverage of dense two-metre-high silver wattle should be 30 to 40 litres of spray per 100m² (10m x 10m) of infestation. For larger areas an equivalent would be 3000 to 4000 litres per infested hectare.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and water courses.

PARTIAL CLEANING (before spraying other labeled or tolerant crops):

After using F.S.A. Clopyralid 600 Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

COMPLETE CLEANING (before spraying susceptible crops):

After using F.S.A. Clopyralid 600 Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate as above, then drain. Quarter fill the tank again and add a liquid alkali detergent (e.g. Surf®, Omo®, Drive®) at 500mL/100L water and circulate throughout the system for at least fifteen minutes. Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using a pressure hose and flush system with clean water.

Plantback Periods: Where F.S.A. Clopyralid 600 Herbicide residue carry over from use rates of less than 250 mL/ha is suspected and susceptible crops are to be planted, test the treated area as follows:

- Field bioassay - where rain allows, plant a small area of the susceptible crop 4 to 6 weeks before desired planting date and take note of any symptoms of injury. If any herbicide symptoms are observed, only plant either canola or a cereal (see recommendation for northern and southern Australia below).
- Pot bioassay - where not practical to do field bioassay, plant a small number of seeds of the susceptible crop into pots containing soil from the treated field. Do this 4 to 6 weeks before desired planting date. If any herbicide symptoms are observed, only plant either canola or cereal (see recommendation for northern and southern Australia below).

Stubble from treated crops - ensure that harvesters effectively spread crop straw and DO NOT leave a heavy 'header trail' after harvest. Burn (if legal in the area), bale and remove, slash or incorporate stubble as soon as practical after harvest and as long as possible before planting next year to allow microbial breakdown of any residues in straw. Heavy stubble loads may carry more residue into the following season. Where heavy stubble burdens and/or non-wetting soils exist and less than recommended rain amount have occurred from application to planting the susceptible crop (see below), only plant a Winter or Summer cereal or canola.

Planting crops following use of F.S.A. Clopyralid 600 Herbicide in previous crop - planting crops 'dry' without significant rain (see below) in the 'Autumn break' increases the risk of injury to susceptible crops. This practice should be avoided, or only plant a Winter or irrigated Summer cereal crop or canola. In severely dry conditions, where less than 30% of average annual rainfall and/or less than the minimum rain (see below) has fallen between application and planting the next year, only plant a Winter or irrigated Summer cereal or canola.

PLANTBACK PERIODS FOR SOUTHERN AUSTRALIA WINTER DOMINANT RAINFALL AREAS (Sth NSW, VIC, SA, WA):

Required rain: A minimum 25 mm rain event in the post-harvest summer to autumn period, with a subsequent extended period of at least 1 week where the top 10 cm of the soil stays moist is required to enable breakdown of soil residues. Fastest residue breakdown will occur under good soil moisture and warm conditions, which promote microbial activity. Where significant rain (>25 mm) has fallen in Summer to Autumn, with soil wetting for at least one week, the following plantback periods apply:

Following Crops	Rate (mL/ha) used previously	Plantback Interval
Clover, chickpea, faba bean, field pea, lentils, lupins, medic and vetch	Up to 150	9 months
	150 - 250	12 months
	>250	24 months
Barley, canola, wheat, oats	All label rates	1 week

PLANTBACK PERIODS FOR NORTHERN AUSTRALIA SUMMER DOMINANT RAINFALL AREAS (Nth NSW, QLD):

Required rainfall before plantback:

If planting susceptible Summer crops - at least 100 mm rain.

If planting susceptible Winter crops - at least 150 mm rain.

This rain or irrigation should wet the soil for extended periods (at least one week) this is essential for breakdown of soil residues prior to planting susceptible crops.

If planting a cereal or canola crop - at least 50 mm of rain or irrigation is required to enable soil wetting for at least one week. Where these requirements have been met the following plantback periods apply:

Following Crops	Rate (mL/ha)	Plantback Interval
	Up to 37.5 mL/ha	37.5 - 150 mL/ha
Chickpea, Cotton, Soybean, Sunflower	3 months	6 months
Lucerne	9 months	9 months
Maize, sorghum	1 week	2 weeks
Barley, canola, wheat, oats	1 week	1 week

Note: Susceptible crops should not be sown for at least 2 years where F.S.A. Clopyralid 600 Herbicide at more than 150 mL/ha has been used in northern Australia.

RESISTANT WEEDS WARNING

GROUP	4	HERBICIDE
--------------	----------	------------------

F.S.A. Clopyralid 600 Herbicide is a member of the Pyridines group of herbicides. The product F.S.A. Clopyralid 600 Herbicide has the disrupters of plant cell growth mode of action. For weed resistance management, the product F.S.A. Clopyralid 600 Herbicide is a Group 4 herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group 4 herbicides. Since occurrence of resistant weeds is difficult to detect prior to use, Four Season Agribusiness Pty Ltd accepts no liability for any losses that may result from

the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or Four Season Agribusiness Pty Ltd representative.

RE-ENTRY PERIODS

Do not allow entry into treated areas until the spray has dried when applying to barley, oats, triticale, wheat, forests (except in control of cape ivy), rights-of-way, industrial situations, and canola, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use. Do not allow entry into treated areas until the spray has dried to undertake low exposure activities (i.e. irrigation, or weeding) and for 8 days after application to undertake high exposure activities (i.e. pruning, training) when applying to control cape ivy in forests, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use. Do not allow entry into treated areas until the spray has dried to undertake low exposure activities (i.e. irrigation) and for 9 days after application to undertake high exposure activities (i.e. hand weeding, transplanting) when applying to pastures and fallow land, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use. Do not allow entry into treated areas until the spray has dried to undertake low exposure activities (i.e. irrigation, scouting or weeding) and for 17 days after application to undertake high exposure activities (i.e. pruning, training) when applying to Pinus radiata plantations, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PRECAUTIONS

When applying to plantation trees including Pinus radiata plantations the following restriction statement applies:

Must use closed mixing and loading for aerial application.

When applying to forests to control groundsel bush, ragwort, silver wattle and cape ivy, the following restriction statement applies: Do not apply by using spraying equipment carried on the back of the user.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures. Composts and mulches - DO NOT apply F.S.A. Clopyralid 600 Herbicide to crops or pastures that will be used for the production of compost or mulches or mushroom substrate. Such compost or mulch made from plant material treated with F.S.A. Clopyralid 600 Herbicide may cause damage to susceptible crops and plants. Susceptible crops and plants include, but are not limited to chickpeas, clover, cotton, faba beans, field peas, fruit trees, lentils, lupins, lucerne, medics, ornamentals, potatoes, safflower, tomatoes, vegetables, grape and kiwifruit vines, vetches, and wattles. Field peas, faba beans, lentils and vetches are particularly susceptible and should not be sown the season following an application of F.S.A. Clopyralid 600 Herbicide at 250 mL/ha.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under WITHHOLDING PERIODS.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

F.S.A. Clopyralid 600 Herbicide has low toxicity to fish, birds, honeybees, livestock, earthworms and aquatic organisms. DO NOT contaminate streams, rivers or watercourses with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seed. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations.

Do not burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above, the wash liquid for disposal.

SAFETY DIRECTIONS

Will irritate the skin. Will damage the eyes. May irritate the nose and throat. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and preparing product for use and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistance gloves, goggles, disposable mist face mask covering mouth and nose. In addition when opening the container and preparing the product for use in aerial spraying equipment wear cotton overalls, over normal clothing, buttoned to the neck and wrist, elbow length chemical resistant gloves and goggles and disposable mist face mask covering mouth and nose. In addition if applying by spraying equipment carried on the back of the user wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat elbow length chemical resistance gloves and goggles and disposable mist face mask covering mouth and nose. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

Causes serious eye damage

**IN A TRANSPORT
EMERGENCY DIAL 000
POLICE OR FIRE BRIGADE**

**FOR SPECIALIST ADVICE IN
AN EMERGENCY ONLY, CALL
1800 033 111 (24 HOURS)**



Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, New South Wales, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255
www.fourseasonsag.com