

READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

OVERSTRIKE



Liquid Sheep Blowfly Treatment

ACTIVE CONSTITUENT: 500 g/L CYROMAZINE



For the control of blowfly strike (*Lucilia cuprina*), which includes organosphosphate resistant strains in long wool sheep for up to fourteen weeks when used as directed.



FOUR SEASONS
AGRIBUSINESS



INDICATIONS

For the control of blowfly strike (*Lucilia cuprina*), which includes organosphosphate resistant strains in long wool sheep for up to fourteen weeks when used as directed. Resistance may develop to any chemical.

DIRECTIONS FOR USE

Restraints

DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

DO NOT collect and re-use fluid which runs off.

Contraindications

This product should not be used on sheep with less than 6 weeks wool as a reduced period of protection may result. Do not use on open wounds, for example, those resulting from marking or mulesing operations.

DOSAGE AND ADMINISTRATION

SHAKE WELL BEFORE USE

| ANIMAL/ PEST | TREATMENT | INITIAL CHARGE | TOPPING UP | CRITICAL COMMENTS |
|--|---|-------------------|-------------------|--|
| Sheep Blowfly (<i>Lucilia cuprina</i>) | Jetting | 2 L per 1000 L | - | <ol style="list-style-type: none">1. Jet at onset of Blowfly wave.2. This product is not recommended for use in jetting races because a reduced period of protection may result.3. Apply 2 to 4 L of jetting fluid per sheep, depending on sheep size and wool length.4. Ensure saturation to the skin.5. Pay particular attention to treatment of the crutch, pizzles and polls. |
| | Dipping (Plunge- Shower and constant replenishment) | 2 L per 1000 L | 2 L per 1000 L | <ol style="list-style-type: none">1. Dip sheep on onset of fly wave.2. Thoroughly saturate sheep. When using a shower dip, run top and bottom sprays independently so as to ensure maximum pressure and therefore penetration.3. The product should not be used alone as a dip for blowfly control of sheep with less than 6 weeks wool as a reduced period of protection may result.4. The product is a non-stripping dip so there is no need to reinforce during dipping. |
| | Dressing | 2 L per 1000 L | - | <ol style="list-style-type: none">1. Clip the wool surrounding the strike, treat the struck area and the surrounding wool with a recommended knock down product then thoroughly jet the sheep with OVERSTRIKE LIQUID.2. The product is a slow killing larvicide and when used on its own, larvae may take 3 to 4 days to die. |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS
AUTHORISED UNDER APPROPRIATE LEGISLATION.**

MIXING

Settling may occur after storage for some weeks. Stir product or invert container several times before opening. Pour the liquid into the vat through a strainer to remove any dry particles or flakes which can occasionally occur under hot storage conditions. Stir well. This product is readily dispersible in both soft and hard water.

DILUTION RATE

Dilution rate: 2 mL per 1 L water

Examples: This means that 400 mL is required to charge a 200 L vat/dip.

MAINTAINANCE OF DIPS

1. Clean out dip, sump and spray lines thoroughly before charging. Disinfect with registered bacteriostat at the manufacturer's recommended rate.
2. DO NOT use old fouled wash. Shower dips should be emptied and cleaned out at the end of each day's dipping. Plunge dips should be emptied and cleaned out within 5 days of charging and more frequently if conditions favour post-dipping lameness infection, except in WA where plunge dips and sumps should be emptied daily.
3. Before use, stir well.

JETTING

Jet the sheep at onset of fly wave with efficient equipment which is able to maintain the pressure of about 550kPa. Use 300-400kPa for lambs and ewes in lamb.

BACTERIOSTAT

This product does not contain a bacteriostat. For control of post dipping infections a registered bacteriostat should be added at the manufacturer's recommended rate.

GENERAL DIRECTIONS

READ DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

This product is designed for use on sheep with at least 6 weeks wool growth (long wool). Dipping sheep heavily infested with grass seeds is not recommended. Heavy rain following the application of this product could diminish the period of protection. If used in jetting races a reduced period of protection may result.

WITHHOLDING PERIOD

MEAT: DO NOT USE less than 7 days before slaughter for human consumption.

WOOL HARVEST INTERVAL: DO NOT USE less than 2 months before shearing or fibre collection.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. Before using this product, confirm the current ESI from Four Seasons Agribusiness Pty Ltd on 1300 449 255 or the APVMA website (www.apvma.gov.au/residues).

Use of this product may result in wool residues that may not comply with European Union Environmental Quality Standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray/dip/dressing wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. When using the prepared spray/dip/dressing wear protective and waterproof clothing and PVC or rubber apron, elbow-length PVC gloves, face shield and water resistant footwear. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

ADDITIONAL USER SAFETY

Additional information is listed in the Safety Data Sheet.

Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre: Phone Australia 131 126.

ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Keep out of reach of children. Store below 30°C (room temperature) in original containers, tightly closed. DO NOT store for prolonged periods in sunlight.

APVMA Approval No.: 91457/132283

Harmful if swallowed. Avoid breathing fumes, mists, vapours or spray. Do not eat, drink or smoke when using this product.

Wear protective gloves, protective clothing and eye or face protection.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF ON SKIN (or hair): Remove immediately all contaminated clothing. Rinse skin with water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.

Continue rinsing. In case of fire, use carbon dioxide, dry chemical, foam, water fog.

**IN AN EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**