



# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE  
OPENING OR USING

F.S.A.

# TRAIL STOP

## Slug and Snail Pellets

ACTIVE CONSTITUENT: 15 g/kg METALDEHYDE

For the control of snails and slugs in many  
agricultural and horticultural systems as  
shown in the Directions For Use

Safety directions and directions for use on  
back of bag



FOUR SEASONS  
AGRIBUSINESS

### SOFTPROOF

Product ID  
**000000**


CLIENT Four Seasons Agribusiness

NAME Trail Stop E-label

FINISHED SIZE 210mm(w) x 297mm(h)

DATE 11th June, 2026

#### COLOURS



APPROVED

NOT APPROVED

DATE: \_\_\_\_\_

SIGNATURE: \_\_\_\_\_

# DGL

#### PLEASE NOTE:

- Colours shown in this proof may vary from the final product compared to what is shown on your screen or what you print out.
- It is the responsibility of the client to ensure all shown information is correct.
- If approved, please indicate above, sign and send back to [labels@dglgroup.com](mailto:labels@dglgroup.com), or you can reply to the email this proof was sent from and confirm approval in writing.
- Some PDF viewing programs can make some text appear thicker than other text. This is a rendering issue within the program and can be resolved by zooming into the artwork.

# TRAIL STOP SLUG AND SNAIL PELLETS

## DIRECTIONS FOR USE

**Restrains:** DO NOT place pellets in heaps.

CROP/SITUATION	PEST	RATE	CRITICAL COMMENTS
Winer cereals, oilseeds	Common white snail ( <i>Ceruella virgata</i> ), White Italian snail ( <i>Theba pisana</i> ), Conical small snail ( <i>Cochlicella acuta</i> ), Grey field slug ( <i>Deroceras reticulatum</i> )	5 – 7.5 kg/ha	Broadcast pellets evenly over the crop or pasture area to be protected from snails and slugs. Pay particular attention to areas adjacent to headlands, fence lines and other areas where snails and slugs shelter and move into crops or pastures. Use the higher rate where there are large populations of snails or slugs and where wet, cool climatic conditions are favourable for snail and slug movement.
Pulses*, Pasture establishment*			<b>*Pulse and pasture crops only:</b> This product is only to be used during the establishment stages of pulse crops and pastures. Broadcast pellets evenly over the crop area to be protected prior to planting, post-plant pre-emergence or early post-emergence (prior to 4 leaf stage).
Around crops and/or border areas where slugs and snails congregate.	Snails, Slugs	50 g/m <sup>2</sup>	Scatter the pellets around the area to be protected and/or where snails and slugs congregate. Spread pellets evenly.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## WITHHOLDING PERIOD

### HARVEST:

**Fruit and vegetables: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.**

**Pulses, Oil seeds, Winter cereals: NOT REQUIRED WHEN USED AS DIRECTED.**

### GRAZING:

**Pastures, Pulses, Oil seeds, Winter Cereals: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 5 WEEKS AFTER APPLICATION.**

## GENERAL INSTRUCTIONS

Scatter pellets using an appropriate applicator. The best time to apply is immediately after rain or during humid, overcast weather when snails and slugs are most active. The pellets eventually decompose leaving no harmful residue. Use regularly to control re-infestation.

### PRECAUTION

DO NOT apply to edible plant parts. DO NOT graze or cut for stockfeed any crops while the pellets are visible in the plant foliage.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, AND ENVIRONMENT

DO NOT contaminate wetlands or watercourses with this product or used containers.

### PROTECTION OF DOMESTIC ANIMALS AND LIVESTOCK

DO NOT place in locations which are accessible to domestic animals, livestock or birds. Keep away from animals, poultry, ducks and domestic pets as accidental poisoning may be fatal.

Dogs and other animals find this bait attractive and it may kill them. If pets or livestock are poisoned contact a Veterinary Surgeon.

### STORAGE AND DISPOSAL

Store in closed, original container in a dry, cool well ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, poultry, seed and fertiliser.

Shake and empty contents into application equipment.

Do not dispose of undiluted chemicals on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and

tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### SAFETY DIRECTIONS

Poisonous if swallowed. Wash hands after use.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

### SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet that is available from the supplier.

### CONDITIONS OF SALE

The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

**IN AN EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**



**FOUR SEASONS**  
AGRIBUSINESS

Four Seasons Agribusiness Pty Ltd  
287 Boorowa Street, Young, New South Wales, Australia, 2594  
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255  
www.fourseasonsag.com