

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

VENGEANCE

pour-on lousicide for sheep

35 g/L IMIDACLOPRID
Also Contains: 300 g/L
N-METHYL-2-PYRROLIDONE (solvent)

For the control of susceptible body lice (*Bovicola ovis*) on shorn sheep up to 7 days offshears and unshorn lambs up to 2 months of age. This product can be expected to protect sheep against with lice for 4 weeks after treatment when applied within 24 hours of shearing.

Vengeance Pour-On is for the control of neonicotinoid susceptible sheep body lice (*Bovicola ovis*) in off-shears (within 7 days of shearing) and unshorn lambs up to 2 months of age.

This product can be used on all adult sheep up to 7 days off-shears and on unshorn lambs up 2 months of age.

Resistance can develop to any chemical.



20L



FOUR SEASONS
AGRIBUSINESS



9 354545 001512

VENGEANCE POUR-ON LOUSICIDE FOR SHEEP

DIRECTIONS FOR USE

Restraint



DO NOT USE on female sheep which are producing or may in future produce milk or milk products for human consumption.

DOSAGE AND ADMINISTRATION

LICE PREVENTION OFF-SHEARS

When using the product up to 7 days off-shears, apply product only with the applicator gun. For sheep over 30kg, apply half the dose as a broadband stripe either side of the midline from the mid-neck to the butt of the tail. The stripes should be no more than 10cm apart. Refer to the diagrams in the Table below. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). DO NOT underdose. Where there is a large variation in size within the group (e.g. lambs), dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

DOSAGE TABLE:

BODYWEIGHT (kg)	DOSE (mL)	METHOD	APPLY IN THE PATTERN BELOW
Below 6 kg	8	Single Stripe	
6.0 - 8.0	12		
8.1 - 10.0	15		
10.1 - 12.5	20		
12.6 - 15	25		
15.1 - 20	30		
20.1 - 30	40		
30.1 - 55	60	Double Stripe	
55.1 - 80	80		
80.1 - 100	90		

Add 10mL for each 10kg above 100kg.

GENERAL DIRECTIONS.

Lice Control

To prevent reinfestation treated sheep must not be mixed with lice free sheep until 6 weeks after treatment. Live lice on ewes treated within 6 weeks of lambing can infect lambs from these ewes. Efficacy is not affected by moderate rainfall either before or after treatment (50 mm rain 1 hour prior to treatment and/or 12.5mm 2 hours after treatment and/or 25 mm 2 hours after treatment). Treated sheep must not be mixed with lousy sheep. Do not use on sheep which have not been clearly shorn for any reason e.g., mycotic dermatitis. Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective live control.

Other Directions

Treated sheep could potentially develop backline dermatitis followed by prolonged exposure to extreme heat and sunlight without access to shade. Do not leave container, hose or applicator in the sun for long periods of time. This product may be used in conjunction with worm control drenches and at the same time as vaccination. This product contains a blue dye to identify treated sheep. The dye is scourable from the wool. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces. Vengeance Pour-On is safe for use on breeding rams.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

WOOL: DO NOT USE less than 2 months before shearing or fibre collection.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 63 before slaughter for export. Before using this product, confirm the current ESI from APVMA website www.apvma.gov.au/residues or by calling Four Seasons Agribusiness on 1300 449 255.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and using the product, wear face shield or goggles. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. If in eyes wash out immediately with water.

ADDITIONAL USER SAFETY

Additional user safety information is listed in the Safety Data Sheet. Rehandling Interval: Do not come into contact with treated sheep until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing).

ENVIRONMENTAL PROTECTION: Imidacloprid is highly toxic to aquatic species. DO NOT contaminate dams, ponds, rivers, waterways or drains with pesticide or used containers.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature). Store in original container, tightly closed in a safe place. Protect from light.

CONDITIONS OF SALE

The sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

APVMA Approval No: 91266/134024

May damage fertility or the unborn child.

May cause respiratory irritation. Causes skin irritation.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

Use only outdoors or in a well-ventilated area.

Avoid breathing mists or spray.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

IF ON SKIN: Wash with plenty of soap and water.

IN AN EMERGENCY, DIAL 000, POLICE OR FIRE BRIGADE

Batch No:

Expiry Date:



FOUR SEASONS
AGRIBUSINESS

Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, New South Wales, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255
www.fourseasonsag.com