

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

SIDEWINDER

POUR-ON FOR CATTLE

ACTIVE CONSTITUENT: 5 g/L MOXIDECTIN

SOLVENT: 150 g/L HYDROCARBON LIQUID

For the treatment and control of Moxidectin sensitive internal and external parasites in cattle and for the treatment and control of lungworm and gastrointestinal roundworms of red deer.



FOUR SEASONS
AGRIBUSINESS

Four Seasons Agribusiness Pty Ltd
287 Boorowa Street, Young, New South Wales, Australia, 2594
ABN 98 115 133 189 ACN 115 133 189 Phone 1300 449 255
www.fourseasonsag.com



INDICATIONS: This product contains Moxidectin, a second-generation macrocyclic lactone endectocide. It is effective in the treatment and control of sensitive strains of the following parasites affecting in **CATTLE:**

Lungworm
Dictyocaulus viviparus

Cattle Tick
Rhipicephalus (Boophilus) microplus

Lice
Linognathus vituli, *Haematopinus eurysternus*,
Solenopotes capillatus, *Bovicola bovis*

Mange Mites
Chorioptes bovis, *Sarcoptes bovis**, *Psoroptes ovis**

*Not found in Australia

Parasite species	Persistent Protection Period
<i>Ostertagia ostertagi</i> , <i>O. lyrata</i>	42 days
<i>Haemonchus placei</i>	28 days
<i>Trichostrongylus axei</i>	28 days
<i>Dictyocaulus viviparus</i>	42 days
<i>Oesophagostomum radiatum</i>	42 days
<i>Bunostomum phlebotomum</i>	42 days
<i>Nematodirus helvetianus</i>	21 days

Buffalo fly control: *Haematobia irritans exigua*
The persistent activity of this product can be used as an aid in control of buffalo fly (*Haematobia irritans exigua*) for up to 14 days after treatment. In addition, this product can be used to treat and control the following parasites of **RED DEER**:

Lungworm
Dictyocaulus viviparus

Parasite species	Persistent Protection Period
<i>Ostertagia</i> spp.	42 days
<i>Oesophagostomum</i> spp.	42 days
<i>Dictyocaulus viviparus</i>	42 days

DOSAGE AND ADMINISTRATION
Cattle and Red Deer: This product should be applied along the top of the back from the withers to the base of the tail.
DOSE RATE: 1 mL per 10 kg liveweight. This will administer the correct dose of 0.5mg/kg liveweight.
 This product can be used in lactating dairy cattle, but **DO NOT RE-TREAT LACTATING CATTLE AT LESS THAN 21 DAYS INTERVALS.**

Cattle heavier than 650 kg should be dosed at 1ml per 10kg liveweight. A representative sample of animals should be weighed before treatment either with scales or weighband. Dose rate should be based on the heaviest animal in each group (bulls, cows, steers, calves, etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.

Product Description: This product comes ready to use as a clear solution containing 5 g/L Moxidectin. When given at a dose volume of 1mL/10 kg liveweight, this product delivers the recommended dosage of 0.5 mg Moxidectin/kg liveweight. This product does not cause irritation to the skin or hide damage when used as directed.

PRODUCT ADVANTAGES
Rainfast: The efficacy of this product is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application. However, the efficacy against tick and buffalo fly may be reduced if very heavy rain occurs following treatment. Treatment of animals under these circumstances is not good agricultural practice.

CONDITIONS OF SALE: the sale, supply, storage, use and application of this product is beyond the control of the manufacturer, and, subject to this provision, all warranties, conditions, rights and remedies express or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application are excluded. Four Seasons Agribusiness Pty Limited and its associated entities shall not accept any liability whatsoever (including consequential loss) or however arising (including negligence) for any damage, injury or death connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

Causes serious eye damage. May cause drowsiness or dizziness. May be fatal if swallowed and enters airways.
Wear protective gloves/protective clothing/eye protection/face protection.
IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician. Do NOT induce vomiting. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. If skin irritation occurs: Get medical advice/attention.

IN AN EMERGENCY, DIAL 000, POLICE OR FIRE BRIGADE

Batch No: Expiry:
